

Belief network inference

Three main approaches to determine posterior distributions in belief networks:

- Exploiting the structure of the network to eliminate (sum out) the non-observed, non-query variables one at a time.
- Search-based approaches that enumerate some of the possible worlds, and estimate posterior probabilities from the worlds generated.
- Stochastic simulation where random cases are generated according to the probability distributions.

A **factor** is a representation of a function from a tuple of random variables into a number.

We will write factor f on variables X_1, \dots, X_j as $f(X_1, \dots, X_j)$.

We can assign some or all of the variables of a factor:

- $f(X_1 = v_1, X_2, \dots, X_j)$, where $v_1 \in \text{dom}(X_1)$, is a factor on X_2, \dots, X_j .
- $f(X_1 = v_1, X_2 = v_2, \dots, X_j = v_j)$ is a number that is the value of f when each X_i has value v_i .

The former is also written as $f(X_1, X_2, \dots, X_j)_{X_1 = v_1}$, etc.

Example factors

$r(X, Y, Z):$

X	Y	Z	val
t	t	t	0.1
t	t	f	0.9
t	f	t	0.2
t	f	f	0.8
f	t	t	0.4
f	t	f	0.6
f	f	t	0.3
f	f	f	0.7

$r(X=t, Y, Z):$

Y	Z	val
t	t	0.1
t	f	0.9
f	t	0.2
f	f	0.8

$r(X=t, Y, Z=f):$

Y	val
t	0.9
f	0.8

$r(X=t, Y=f, Z=f) = 0.8$

Multiplying factors

The **product** of factor $f_1(\bar{X}, \bar{Y})$ and $f_2(\bar{Y}, \bar{Z})$, where \bar{Y} are the variables in common, is the factor $(f_1 \times f_2)(\bar{X}, \bar{Y}, \bar{Z})$ defined by:

$$(f_1 \times f_2)(\bar{X}, \bar{Y}, \bar{Z}) = f_1(\bar{X}, \bar{Y})f_2(\bar{Y}, \bar{Z}).$$

Multiplying factors example

f_1 :

A	B	val
t	t	0.1
t	f	0.9
f	t	0.2
f	f	0.8

f_2 :

B	C	val
t	t	0.3
t	f	0.7
f	t	0.6
f	f	0.4

$f_1 \times f_2$:

A	B	C	val
t	t	t	0.03
t	t	f	0.07
t	f	t	0.54
t	f	f	0.36
f	t	t	0.06
f	t	f	0.14
f	f	t	0.48
f	f	f	0.32

Summing out variables

We can **sum out** a variable, say X_1 with domain $\{v_1, \dots, v_k\}$, from factor $f(X_1, \dots, X_j)$, resulting in a factor on X_2, \dots, X_j defined by:

$$\begin{aligned} & \left(\sum_{X_1} f \right) (X_2, \dots, X_j) \\ &= f(X_1 = v_1, \dots, X_j) + \dots + f(X_1 = v_k, \dots, X_j) \end{aligned}$$

Summing out a variable example

f_3 :

A	B	C	val
t	t	t	0.03
t	t	f	0.07
t	f	t	0.54
t	f	f	0.36
f	t	t	0.06
f	t	f	0.14
f	f	t	0.48
f	f	f	0.32

$\sum_B f_3$:

A	C	val
t	t	0.57
t	f	0.43
f	t	0.54
f	f	0.46

If we want to compute the posterior probability of Z given evidence $Y_1 = v_1 \wedge \dots \wedge Y_j = v_j$:

$$\begin{aligned} P(Z|Y_1 = v_1, \dots, Y_j = v_j) & \\ &= \frac{P(Z, Y_1 = v_1, \dots, Y_j = v_j)}{P(Y_1 = v_1, \dots, Y_j = v_j)} \\ &= \frac{P(Z, Y_1 = v_1, \dots, Y_j = v_j)}{\sum_Z P(Z, Y_1 = v_1, \dots, Y_j = v_j)}. \end{aligned}$$

So the computation reduces to the probability of $P(Z, Y_1 = v_1, \dots, Y_j = v_j)$.

We normalize at the end.

Probability of a conjunction

Suppose the variables of the belief network are X_1, \dots, X_n .
To compute $P(Z, Y_1 = v_1, \dots, Y_j = v_j)$, we sum out the other variables, $Z_1, \dots, Z_k = \{X_1, \dots, X_n\} - \{Z\} - \{Y_1, \dots, Y_j\}$.
We order the Z_i into an **elimination ordering**.

$$\begin{aligned} &P(Z, Y_1 = v_1, \dots, Y_j = v_j) \\ &= \sum_{Z_k} \cdots \sum_{Z_1} P(X_1, \dots, X_n)_{Y_1 = v_1, \dots, Y_j = v_j} \\ &= \sum_{Z_k} \cdots \sum_{Z_1} \prod_{i=1}^n P(X_i | \text{parents}(X_i))_{Y_1 = v_1, \dots, Y_j = v_j} \end{aligned}$$

Computing sums of products

Computation in belief networks reduces to computing the sums of products.

- How can we compute $ab + ac$ efficiently?

Computing sums of products

Computation in belief networks reduces to computing the sums of products.

- How can we compute $ab + ac$ efficiently?
- Distribute out the a giving $a(b + c)$

Computing sums of products

Computation in belief networks reduces to computing the sums of products.

- How can we compute $ab + ac$ efficiently?
- Distribute out the a giving $a(b + c)$
- How can we compute $\sum_{Z_1} \prod_{i=1}^n P(X_i | \text{parents}(X_i))$ efficiently?

Computing sums of products

Computation in belief networks reduces to computing the sums of products.

- How can we compute $ab + ac$ efficiently?
- Distribute out the a giving $a(b + c)$
- How can we compute $\sum_{Z_1} \prod_{i=1}^n P(X_i | \text{parents}(X_i))$ efficiently?
- Distribute out those factors that don't involve Z_1 .

Variable elimination algorithm

To compute $P(Z|Y_1 = v_1 \wedge \dots \wedge Y_j = v_j)$:

- Construct a factor for each conditional probability.
- Set the observed variables to their observed values.
- Sum out each of the other variables (the $\{Z_1, \dots, Z_k\}$) according to some elimination ordering.
- Multiply the remaining factors. Normalize by dividing the resulting factor $f(Z)$ by $\sum_Z f(Z)$.

Summing out a variable

To sum out a variable Z_j from a product f_1, \dots, f_k of factors:


- Partition the factors into
 - ▶ those that don't contain Z_j , say f_1, \dots, f_i ,
 - ▶ those that contain Z_j , say f_{i+1}, \dots, f_k

We know:

$$\sum_{Z_j} f_1 \times \dots \times f_k = f_1 \times \dots \times f_i \times \left(\sum_{Z_j} f_{i+1} \times \dots \times f_k \right).$$

- Explicitly construct a representation of the rightmost factor. Replace the factors f_{i+1}, \dots, f_k by the new factor.

Variable elimination example



$$\left. \begin{array}{l} P(A) \\ P(B|A) \end{array} \right\} \xrightarrow{\text{elim } A} f_1(B)$$

$$\left. \begin{array}{l} P(C) \\ P(D|BC) \\ P(E|C) \end{array} \right\} \xrightarrow{\text{elim } C} f_2(BDE)$$

$$\begin{array}{l} P(F|D) \\ P(G|FE) \end{array}$$

$$P(H|G) \left. \right\} \xrightarrow{\text{obs } H} f_3(G)$$

$$P(I|G) \left. \right\} \xrightarrow{\text{elim } I} f_4(G)$$