

Solutions to Homework 1

Exercise 1

Adding page 5 as asked (pages 3 and 5 link to each other) gives the new link matrix:

$$A = \begin{bmatrix} 0 & 0 & \frac{1}{2} & \frac{1}{2} & 0 \\ \frac{1}{3} & 0 & 0 & 0 & 0 \\ \frac{1}{3} & \frac{1}{2} & 0 & \frac{1}{2} & 1 \\ \frac{1}{3} & \frac{1}{2} & 0 & 0 & 0 \\ 0 & 0 & \frac{1}{2} & 0 & 0 \end{bmatrix}$$

As before, we are looking for an eigenvector $\mathbf{x} = [x_1, x_2, x_3, x_4, x_5]^T$ corresponding to the eigenvalue 1 (which exists because $\mathbf{A}$ is column-stochastic). This $\mathbf{x}$ will be our ranking. Solve $\mathbf{Ax} = \mathbf{x}$, i.e. $(\mathbf{A} - \mathbf{I})\mathbf{x} = 0$, for $\mathbf{x}$:

$$-x_1 + \frac{1}{2}x_3 + \frac{1}{2}x_4 = 0 \quad (1)$$

$$\frac{1}{3}x_1 - x_2 = 0 \quad (2)$$

$$\frac{1}{3}x_1 + \frac{1}{2}x_2 - x_3 + \frac{1}{2}x_4 + x_5 = 0 \quad (3)$$

$$\frac{1}{3}x_1 + \frac{1}{2}x_2 - x_4 = 0 \quad (4)$$

$$\frac{1}{2}x_3 - x_5 = 0 \quad (5)$$


Solve ‘by hand’ first, then confirm solution with MATLAB. Set $x_3 = 2$ and $x_5 = 1$ to satisfy equation (5). Then subtract (4) from (3) to obtain $x_4 = 2/3$, plug x_3 and x_4 into equation (1) to get $x_1 = 4/3$. Finally plug x_1 into equation (2) to get $x_2 = 4/9$.

So we’ve got one eigenvector: $\mathbf{x} = [4/3, 4/9, 2, 2/3, 1]^T$ corresponding to $\lambda = 1$; there are many more solutions - as many as you’d like, all collinear to $\mathbf{x}$. (We can agree to pick the one that ensures the components of $\mathbf{x}$ sum to 1, i.e. scale $\mathbf{x}$ by dividing each component by the sum of $\mathbf{x}$ ’s components.)

Here's some MATLAB that does the same:

```
A = [ 0      0      1/2    1/2     0 ;
      1/3    0      0      0      0 ;
      1/3    1/2    0      1/2    1 ;
      1/3    1/2    0      0      0 ;
      0      0      1/2    0      0 ];

[V D] = eig(A1);
disp('the eigenvector associated with eigenvalue 1:')
V(:,1)
x = V(:,1)/sum(V(:,1))
figure(1)
bar(x)
```



Thus the new ranking is

solution

$$x = [0.2449, 0.0816, 0.3673, 0.1224, 0.1837]$$

and page 3 is now ranked the highest.

Exercise 7

Let $\mathbf{A}$ and $\mathbf{S}$ be column stochastic matrices, i.e. $\sum_i A_{ij} = 1$ and $\sum_i S_{ij} = 1$ for every j -th column of $\mathbf{A}$ or $\mathbf{S}$. Let $0 \leq m \leq 1$ and let $\mathbf{M} = m\mathbf{A} + (1 - m)\mathbf{S}$. Consider the sum of the elements in $\mathbf{M}$'s j -th column:

$$\begin{aligned} \sum_i M_{ij} &= \sum_i (mA_{ij} + (1 - m)S_{ij}) \\ &= \sum_i mA_{ij} + \sum_i (1 - m)S_{ij} \\ &= m \sum_i A_{ij} + (1 - m) \sum_i S_{ij} \\ &= m + (1 - m) \\ &= 1 \end{aligned}$$

This being true for any j , $\mathbf{M}$ is column-stochastic.

Exercise 11

Here the link matrix $\mathbf{A}$ is the same as in problem 1. Set $m = 0.15$ and write down the matrix $\mathbf{M} = 0.85\mathbf{A} + 0.15\mathbf{S}$ where $\mathbf{S}$ is the 5×5 matrix each element of which is $1/5$.

$$\mathbf{A} = \begin{bmatrix} 0 & 0 & 1/2 & 1/2 & 0 \\ 1/3 & 0 & 0 & 0 & 0 \\ 1/3 & 1/2 & 0 & 1/2 & 1 \\ 1/3 & 1/2 & 0 & 0 & 0 \\ 0 & 0 & 1/2 & 0 & 0 \end{bmatrix};$$

```
S = 1/5 * ones(5);
m = 0.15;
M = (1 - m)*A + m*S
[V D] = eig(M)
% diag(D)
ranks = V(:,1) / sum(V(:,1))
bar(ranks)
```

Here the output is:

~~ranks~~ = [0.2371, 0.0972, 0.3489, 0.1385, 0.1783]

Slightly different scores but the relative ranking of the pages is the same.

solutions

page 3

